

Shonihari

Week 6 (4) 2/18

Focus on five phase pattern based on comparative pulse dx and pulse quality

- ◆ Practice root treatment diagnosis and treatment (tonification and dispersion) with Teishin.

TIPS FOR SHONIHARI

Perform a pulse dx. in order to determine the SHO.

Characteristic kid's calm Pulse quality

Floating, Fast, Soft, Short

Order of Pulse Diagnosis

Is it Yin or Yang disease

- Yang disease; It's from OPI(Jaki)
- Yin disease; It's constitutional or It's influenced by parent EPI.
- Catch the middle pulse then find excess

@ Yang disease

Acute: Excess on yang meridian with yin deficiency in same phases.

Fetal toxin/Chronic: Blood stasis

- ★ LR ex. w/ LU+KD def. Many of illness
- ★ LR ex. w/ SP def. Food Allergy
- ★ LU ex. w/ SP def. Allergy/Eczema/Asthma

@ Yin disease

Yin Ouki-Jitsu

(Excess without Jaki) with Yin deficient on control cycle

- ★ After tonify primary dif. Yin then apply the **Yusha/Shanting**/Tonification on the control cycle point on Excess meridian.

Treatment Frequency

1. General or

Kan-no-mushi (irascibility bags syndrome)

On average _____ every day for 4-5 days.

In light cases _____ every day for 2-3 days.

In serious cases _____ every day for 7-10 days.

2. Judging the effectiveness of treatment

Questioning the guardian in detail. If the symptom worsens, the amount of stimulation should be decreased to 1/5 - 1/10 of the original dosage.

3. Timing of the end of treatment


You must end it, when the normal body is approached most.

How to use Teishin for dispersion

Use round tip for weak constitution or Deficient type of Ex. Jaki

Sharp tip for general Excess type of Jaki excess and /or very rough skin (blood stasis).

Get the Picture



Have More to Say?

Most effective point for Ac-tx. is Volcano point. For Local Tx. Popular Basic Point Selection for Root Tx.

@Yin

Lung shō - LU-9, SP-3

Spleen shō - PC-7, SP-3

Liver shō - LV-8, KI-10

Kidney shō - LU-5 or LU-8 and KI-7

@Yang

Luo point